Things Parents
Want to Know
About Newborn
Screening From
Their Child's
Health Professional








141 Northwest Point Boulevard Elk Grove Village, Illinois 60007 847/434-4000

www.aap.org



Newborn Screening Disorders

What parents want to know about newborn screenings disorders.

Quick Reference to Newborn Screening Disorders

7 Things Parents Want to Know About Newborn Screening From Their Child's Health Professional

Biotinidase Deficiency (BIO)

BIO is a enzyme deficiency that occurs in about 1 in 60,000 newborns and can result in seizures, hearing loss and death in severe cases. Treatment is simple and involves daily doses of biotin.

Congenital Adrenal Hyperplasia (CAH)

CAH is caused by decreased or absent production of certain adrenal hormones. The most prevalent type is detected by newborn screening in about 1 in 15,000 newborns. Early detection can prevent death in boys and girls and sex misassignment in girls. Treatment involves lifelong hormone replacement therapy.

Congenital Hypothyroidism (CH)

Inadequate or absent production of thyroid hormone results in CH and is present in about 1 in 3,500 newborns. Thyroid hormone replacement therapy begun by 1 month of age can prevent mental and growth retardation.

Congenital Toxoplasmosis (TOXO)

Infection of the fetus with a parasite ingested by the mother during pregnancy can result in TOXO in the newborn. The transmission rate is about 30% and the national incidence is approximately 1 in 10,000 newborns based on limited screening. Early diagnosis and drug therapy reduces the risk of blindness, mental retardation or other serious complications.

Cystic Fibrosis (CF)

CF occurs in about 1 in 4,000 U.S. newborns and is characterized by progressive lung disease, pancreatic dysfunction and other organ failures. Confirmatory testing usually involves sweat testing and treatment leads to decreased hospitalizations and better nutritional and pulmonary outcomes. Caucasians are at an increased risk.

Galactosemia (GAL)

Failure to metabolize the milk sugar galactose results in GAL and occurs in about 1 in 50,000 newborns. The classical form detected by newborn screening can lead to cataracts, liver cirrhosis, mental retardation and/or death. Treatment is elimination of galactose from the diet usually by substituting soy.

Homocystinuria (HCY)

HCY is caused by an enzyme deficiency that blocks the metabolism of an amino acid that can lead to mental retardation, osteoporosis and other problems if left undetected and untreated. The incidence is approximately 1 in 350,000 U.S. newborns. Treatment may involve dietary restrictions and supplemental medicines.

Maple Syrup Urine Disease (MSUD)

MSUD is a defect in the way that the body metabolizes certain amino acids and is present in about 1 in 200,000 U.S. newborns. Early detection and treatment with dietary restrictions can prevent death and severe mental retardation. There is an increased risk in Mennonites.

Medium Chain Acyl-CoA Dehydrogenase (MCAD) Deficiency

The most common disorder in the way the body metabolizes fatty acids is called MCAD deficiency. Undetected, it can cause sudden death. Treatment is simple and includes ensuring regular food intake. The incidence from newborn screening is not yet known, but is thought to be approximately 1 in 15,000 newborns.

Phenylketonuria (PKU)

An enzyme defect that prevents metabolism of phenylalanine, an amino acid essential to brain development, is known as PKU and occurs in approximately 1 in every 19,000 U.S. newborns. Undetected and untreated with a special diet, PKU leads to irreversible mental retardation. Persons of European descent are at increased risk.

Sickle Cell Disease (SCD)

Sickle cell anemia (Hb-SS-Disease) is the most prevalent SCD and causes clogged blood vessels resulting in severe pain and other severe health problems. Other common SCDs include Hb-SC-Disease and various thalassemias. Newborn screening detects about 1 in 2,500 newborns with SCD annually. Persons of African or Mediterranean descent are at an increased risk.

Other Fatty Acid Oxidation (FAO) Disorders

Besides MCAD deficiency, other FAO disorders may be detected through newborn screening. They are usually described in categories based on the length of the fatty acid involved. Undetected and untreated they can cause seizures, coma and even death. The incidences of various FAO disorders are not known since it is only recently that early detection through newborn screening has occurred.

Organic Acid (OA) Disorders

Organic acidemias are a group of metabolic disorders that lead to accumulation of organic acids in the blood and urine and may be detected in newborn screening though analysis of acylcarnitine profiles. Symptoms can be diminished by restricting protein in the diet and supplementation with vitamins and/or carnitine. Because newborn screening for these disorders is relatively new, the degree of occurrence in newborns is not yet known.

Urea Cycle Disorders (UCD)

A UCD is a genetic disorder caused by a deficiency of one of the enzymes responsible for removing ammonia from the blood stream. Some UCDs may be detected as a part of newborn screening. They are characterized by seizures, poor muscle tone, respiratory distress, and coma, and result in death if left undetected and untreated. Because newborn screening for these disorders is relatively new, the degree of occurrence in newborns is not yet known.

The Health Professional's Guide for Brief Prenatal Discussion with Parents

- 1) All newborn babies are required by the State to get tested for some rare disorders before they leave the hospital.
- 2) Babies with these disorders may look healthy at birth.
- 3) Serious problems can be prevented if we find out about the disorders right away.
- 4) To do the test, a nurse will take a few drops of blood from your baby's heel.
- 5) Your baby's health professional and the hospital will get a copy of the test results. Call your baby's health professional if you would like to talk about the results.
- 6) Some babies will need to be retested. If your baby needs to be retested, you will be notified. It is very important to get retested quickly.
- 7) If you have further questions you can call the clinic or the State Health Department.

